



Band structure of an ionic solid: The case of MgO



Objectives

- Compute and analyze the band structure of an ionic solid

MgO an ionic solid that crystallizes in the rocksalt structure

Go to the directory where the exercise of the bands of MgO is included
Inspect the input file, MgO.fdf

```
SystemName Magnesium Oxide Crystal
SystemLabel      MgO

NumberOfAtoms      2
NumberOfSpecies     2

%block Chemical_Species_Label
  1  12  Mg
  2   8   O
%endblock Chemical_Species_Label

LatticeConstant    4.117 Ang
%block LatticeVectors
  0.000  0.500  0.500
  0.500  0.000  0.500
  0.500  0.500  0.000
%endblock LatticeVectors

AtomicCoordinatesFormat Fractional
%block AtomicCoordinatesAndAtomicSpecies
  0.000  0.000  0.000  1
  0.500  0.500  0.500  2
%endblock AtomicCoordinatesAndAtomicSpecies

%block kgrid_Monkhorst_Pack
  6  0  0  0.5
  0  6  0  0.5
  0  0  6  0.5
%endblock kgrid_Monkhorst_Pack
```

More information at the [Siesta web page](http://www.icmab.es/siesta)
<http://www.icmab.es/siesta> and follow the
link Documentations, Manual

The equilibrium lattice constant within
LDA has been computed for you...

Rocksalt structure:


FCC lattice

+ a basis of two atoms

Sampling in k in the first Brillouin zone
to achieve self-consistency

Once SCF has been achieved, we compute the bands along the high symmetry points in the First-Brillouin zone

First-Brillouin zone of a FCC ,
with the high symmetry points



New variables to plot the band structure

BandLinesScale				pi/a	
%block BandLines					
1	1.5	1.5	0.0	K	# Begin at K
38	0.0	0.0	0.0	\Gamma	# 38 points from K to Gamma
36	0.0	2.0	0.0	X	# 36 points from Gamma to X
18	1.0	2.0	0.0	W	# 18 points from X to W
26	1.0	1.0	1.0	L	# 26 points from W to L
31	0.0	0.0	0.0	\Gamma	# 31 points from L to Gamma
%endblock BandLines					

Once SCF has been achieved, we compute the bands along the high symmetry points in the First-Brillouin zone

Check that you have all the required files

A pseudopotential file (.vps or .psf) for every atomic specie included in the input file

For Mg and O within LDA, you can download it from the Siesta web page.

Run the code,

```
siesta < MgO.fdf > MgO.out
```

Wait for a few seconds... and then you should have an output, and a file called **MgO.bands**

If you inspect this file, you will find something like

```
-5.01887068217278  
0.0000000000000000E+000 3.33847877261664  
-25.1877901161284 143.069317296392  
18 1 150  
0.000000 -23.1506 -11.6755 -10.7473 -9.0169 4.0923 ...  
...
```

Coordinate of the k-point in the path, and eigenvalues (in eV).
There are as many eigenvalues as orbitals in the unit cell.

Energy of the Fermi level

Minimum and maximum length of the path in k-space

Minimum and maximum eigenvalues

Number of orbitals in the unit cell, number of different spin polarization, and number of k-points in the walk through the 1BZ

Once SCF has been achieved, we compute the bands along the high symmetry points in the First-Brillouin zone

To plot the band structure, there is a Utility in the directory Util, called gnumbands.f

To use it:

```
cp ~/siesta/Util/gnumbands.f .
```

```
<your_fortran_compiler> -o gnumbands.x gnumbands.f
```

```
gnumbands.x < MgO.bands > MgO.bands.gnuplot.dat
```

The name of this file is free

```
gnuplot
```


```
plot "MgO.bands.gnuplot.dat" using 1:2 with lines
```

```
set xrange [0:3.34] → 3.34 is the position of the last k-point in the path in k-space
```


```
set yrange [-30.0:25.0] → this is large enough to include all the valence bands
```

```
replot
```

Once SCF has been achieved, we compute the bands along the high symmetry points in the First-Brillouin zone



The most important point: analyze your results



The Fermi energy lies in a gap \Rightarrow insulator

Theo. direct gap = 5.3 eV

Expt. Gap = 7.8 eV

(LDA band gap underestimation)

The most important point: analyze your results

H																	He
Li	Be											B	C	N	O	F	Ne
Na	Mg											Al	Si	P	S	Cl	Ar
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
Cs	Ba		Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
Fr	Ra		Rf	Db	Sg	Bh	Hs	Mt	Uun	Uuu	Uub						
		La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	
		Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr	




Mg loses two electrons
that are gained by O



One would expect O (one *s* band and three *p* bands) bands completely full
and Mg bands completely empty

The most important point: analyze your results



The Fermi energy lies in a gap \Rightarrow insulator

Theo. direct gap = 5.3 eV

Expt. Gap = 7.8 eV


(LDA band gap underestimation)

O 2p

(three bands)

O 2s


Once the transfer of charge is produced,
the atoms only interact electrostatically



In a very simplified tight-binding model,
the width of the band is proportional to the interactions with nearest neighbours

If the interaction is small, the bands are very flat.

The most important point: analyze your results



The Fermi energy lies in a gap \Rightarrow insulator

Theo. direct gap = 5.3 eV

Expt. Gap = 7.8 eV

(LDA band gap underestimation)

O 2p

(three bands)

O 2s

Very small dispersion of the O s and O p bands